

E03: Hyperkalemia

Andrew Mills

Updated: December 07, 2020

Reviewed:

Introduction

Although there are many potential electrolyte disturbances, hyperkalemia is arguably the most serious. In addition, it may be reasonably identified and treated in the pre-hospital environment based on clinical features. The strict laboratory testing diagnosis of hyperkalemia is a serum potassium level over 5.5 mmol/L.

In rare cases with signs of hemodynamic compromise and potentially life-threatening arrhythmias, a clinical suspicion of hyperkalemia may be sufficient for initiating treatment.

Essentials

- The lethality of hyperkalemia is directly related to the rapidity with which the condition has developed, in addition to the absolute level of serum potassium.
- Correlation of specific ECG changes with specific serum levels has not been adequately demonstrated.
- Clinical suspicion of hyperkalemia alone is not cause for treatment in the prehospital setting.
- Treatment of life-threatening hyperkalemia aims at preventing or resolving lethal arrhythmias and restoring hemodynamic stability; this can be accomplished by stabilizing the myocardium, shifting potassium back into the intracellular space, and removing excess potassium from the body. The majority of these interventions are only available in hospital.

Additional Treatment Information

- Bradyarrhythmias with bizarre morphologies should prompt a strong consideration of hyperkalemia.
- To warrant prehospital intervention, patients must present with significant hemodynamic or arrhythmogenic instability, alongside a suspicion of hyperkalemia as the likely cause.

General Information

- Classic causes of hyperkalemia:
 - Increased intake, either through potassium supplementation or diet
 - Increased production, as occurs in hemolysis, rhabdomyolysis, extensive burns, or as a result of intense physical activity or trauma (particularly crush injuries and tissue ischemia)
 - Decreased excretion, caused by acute or end-stage chronic renal failure, or by some drugs (such as nonsteroidal anti-inflammatory drugs, cyclosporine, potassium-sparing diuretics, and ACE inhibitors)
 - Shifts from intracellular to extracellular fluid as a result of acidosis (either metabolic or respiratory), insulin deficiency, or some drugs (particularly succinylcholine in certain populations, beta blockers, and digoxin)
- Clinical features of hyperkalemia are often non-specific:
 - Generalised muscle weakness, paresthesia and/or absent deep tendon reflexes. In rare cases, muscular paralysis and hypoventilation may be observed.
 - Mental status change including confusion, fatigue and lethargy
 - Signs of renal failure, such as edema, skin changes, and dialysis sites, may be present
- The ECG is one of the most important diagnostic tools in detecting hyperkalemia. ECG changes associated with Hyperkalemia include:
 - Tall tented T-waves
 - Flattened or absent P waves
 - Prolonged PR Interval
 - Wide QRS
 - Bradycardia
- These changes may progress to bizarre QRS complexes, sine waves, or asystole.

Interventions

First Responder

- Keep patient at rest in a position of comfort.
- Oxygen as required to maintain SpO₂ ≥ 94%
- → [A07: Oxygen and Medication Administration](#)
- Rapid transport
- Consider ACP intercept

Advanced Care Paramedic – All FR, EMR, and PCP interventions, plus:

- Obtain vascular access.
- → [D03: Vascular Access](#)
- In patients with significant hemodynamic instability or dysrhythmia, and a suspicion of hyperkalemia:
 - **REQUIRES CLINICAL CONSULTATION (1-833-829-4099)**
 - Stabilize cellular action potential:
 - [Calcium chloride](#). May repeat after 5 minutes if ECG changes persist or recur.
 - Shift potassium intracellularly:
 - [Sodium bicarbonate](#)
 - [Salbutamol](#)

Critical Care Paramedic – All FR, EMR, PCP, and ACP interventions, plus:

- Stabilize cellular action potential:
 - Calcium gluconate IV: 1.0 g slow push over 2-3 minutes. May repeat once after 5 minutes if ECG changes persist or recur.
- Shift potassium intracellularly:
 - D10W with 10-20 U insulin R mixed: give 500 mL intravenously over 60 minutes, or:
 - Insulin R 10 units IV followed by glucose 25 g IV.
 - [Sodium bicarbonate](#) IV: 150 mEq in 1 L D5W over 2-4 hours depending on volume status.
- Eliminate potassium:
 - Furosemide IV: 40 mg every 12 hours

References

1. Mount DB. Treatment and prevention of hyperkalemia in adults. In UpToDate. 2019. [[Link](#)]
2. Lehnhardt A et al. Pathogenesis, diagnosis and management of hyperkalemia. 2011. [[Link](#)]
3. North Carolina Office of EMS. Dialysis/Renal Failure. 2012. [[Link](#)]
4. Porter RS et al. The Merck manual of diagnosis and therapy. 20th edition. 2018.

